Table 1.—Solar radiation intensities during April, 1927

## Washington, D. C.

[Gram-calories per minute per square centimeter of normal surface]

				£	Sun's z	enith o	listanc	е			
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noor
Date	75th mer.				A	ir mas	38				Loca
	time		Α.	м.				Ρ,	М,		solar time
	е.	5.0	4.0	3.0	2.0	1 1.0	2.0	3.0	4.0	5.0	е.
Apr. 6	mm. 6.27	cal.	cal.	cal.	cal.	cal.	cal. 1.02	cal. 0.68	cal. 0.57	cal.	mm. 6,50
Apr. 7 Apr. 11 Apr. 12	3.63 2.62 4.75	0, 73 0, 86 0, 85	0. 86 1. 01 0. 89	1.11	1.32		1.15 1.29	1. 09			4.17
Apr. 14	2.87 14.10	0.77	0.92	1.14	1.30	1.29	0.87	1.08			3. 63 12. 68 3. 30
Apr. 25 Apr. 28		0. 67 0. 66		0.98		1.49	1. 19	l			3. 99 4. 57
Means Departures		0, 76 +0, 05			1. 22 +0. 14					(0. 86) +0. 24	
				Madi	son, V	/is.					
Apr. 5	5. 16				1.30						5. 56 5. 16
Apr. 6 Apr. 20 Apr. 23	3, 15 3, 15			1.14		1.47					3. 18 2. 06 6. 50
Apr. 27 Apr. 30					1, 11 1, 22	1.31 1.45	1.18				5. 16
Means Departures				(1. 14) +0. 05	1. 24 +0, 02						
<del></del>	<u> </u>	·	<u>.                                    </u>	<u>'</u>	In, Ne	<u></u>		`	•	<u>'</u>	

	Lincoln, Nebr.										
Apr. 4	8.18		<u> </u>		 	1. 55	1.38	1. 23	1.06	0.92	
Apr. 6 Apr. 17	5.36 7.57		0.88	1.04	1.13 1.22						5. 16 7. 04
Apr. 18	8.48			0. 90			1.32	1, 15	1.00		13. 13 5. 56
Apr. 19 Apr. 21	11.38 2.87		1. 05								2. 26
Apr. 22 Apr. 25	3.30 6.27		0.93								3.00 8.18
Apr. 26	7.87 5.79		0, 60	0. 91 0. 83			1. 21		<b></b> -		6. 27 6. 27
Apr. 27 Apr. 29	7. 87			0.00		1. 47	1. 16	0, 93	0.79	0.69	
Means		(0. 81)	0. 87	1. 00	1. 20	1. 50	1. 27			(0. 80)	   <b>-</b>
Departures		+0.07	+0.04	+0. 02	-0.01	+0.05	+0.09	+0.12	+0.11	+0.09	
	<u> </u>	<u>'</u>		·	<u> </u>		<u>'</u>	<u>'</u>	·	·	<u>'</u>

<sup>1</sup> Extrapolated.

Table 2.—Solar and sky radiation received on a horizontal surface

[Gram-calories per square centimeter of horizontal surface]

Week be-	Average daily radiation from								daily departure m normal		
ginning—	Wash- ington	Madi- son	Lincoln	Chi- cago	New York	Twin Falls	Wash- ington	Madi- son	Lincoln		
1927 Apr. 2 Apr. 9 Apr. 16 Apr. 23	259 430 402 404	315 373 326 378	344 155 438 485	383 284 271 333	283 423 278 373	497 467 536 648	-122 +19 -25 -24	-67 -28 -78 -51	-73 -264 -10 +17		
Deficiency	since fir	st of year	r on Apr.	29			-4, 830	-2, 975	-5, 698		

## POSITIONS AND AREAS OF SUN SPOTS

[Communicated by Capt. Edwin T. Pollock, Superintendent U. S. Naval Observatory]
[Data furnished by Naval Observatory, in cooperation with Harvard, Yerkes, and Mount Wilson observatories]

	East	ern	Heliog	raphic	Area 1		
Date	stand civil t	lard	Longi- tude	Latitude	Spot	Group	
1927	h.	m.	0	•			
far. 9 (Yerkes)	12 12	10 40	+6.9 $-62.3$	14.5 +16.0	73 97		
pr. 3 (Naval Observatory)	14	3	-56.0 -43.0	+14.0 +13.5	93		
	ĺ		-38.0	-24.0	62	71	
			-10.5 -8.5	-15.0	46	1	
		1	-4. 0	+17.5 -13.0			
	1	- 1	$+54.0 \\ +61.5$	+15.5 -22.5		34	
pr. 4 (Mount Wilson)	14	25	-41.0	+14.0	97		
•			-41. 0	+10.0	1	72	
			-27.5 $-25.0$	+12.5 $-24.5$	20		
			-10.0	-16.0	71		
			+6.0 +7.0	+16.0 -8.0			
			+7.0	-14.0			
		1	+12.0 +46.0	+22.0 -14.0		1	
			<del>+</del> 70. 0	+14.5	3		
pr. 5 (Yerkes)	9	56	+73.0 -80.0	-23. 0 -15. 5	485	4	
pri o (1 oraco)	•	-	<b>30</b> . 5	+13.5	75		
			-24.0 -19.0	+15.0 +13.0	75 100		
			-15.0	+11.0	120		
	Į		+16.0 +25.0	+16.0 +22.0	95 50		
			+79.0	-29.0	480		
pr. 6 (Naval Observatory)	11	46	-82. 0 -69. 0	-17. 0 -13. 5		2	
			-16.0	+14.5	62		
			-5.0 +30.0	+14.0 +16.5	42	4	
			+37.0	+22.0			
pr. 7 (Naval Observatory)	11	47	+71.0 -68.0	-13. 5 -17. 0		1 2	
pit i (ivavai Observatory)		Ξ'	-55.0	-14.0		2	
			+7.0 +42.5	+12.5 +16.5	31	4	
	1	-	+50.0	+22.0			
pr. 10 (Naval Observatory)	12	7	+85.0 -79.0	-15.0 +30.0	123 216		
pr. 10 (14avar Obsorvatory)	1.2	•	-78.0	-7.0		2	
			-45.0 -38.5	-22.5 +13.5		1	
			-26.0	-17.5		2	
			-14.0 +34.5	-14.0 +13.5		1	
			+51.0	+12.5		4	
pr. 11 (Naval Observatory)	11	51	+72.0 -80.0	+5.0 +11.0		3	
pr. 11 (tyavar Observatory)	11	31	72.0	+33.0		7	
		- 1	-69.0 -68.0	-7. 0 -14. 5			
			-60. 5	<b>—7.</b> 5			
			-29.0	-21.5		١,	
			-25.0 -20.0	+14.0 -18.5	9	1	
	i		-10.5	-18.0		2	
	İ		-0.5 +63.0	-13.0 +13.0		i a	
pr. 12 (Naval Observatory)	11	46	<b>-67.0</b>	+13.0 +11.5		. 2	
			-58. 0 -57. 0	+32.5 -5.5	25	2	
			-53.0	-15.0		. 1	
			-47. 5 -15. 0	-7.5 $-21.5$		1	
			<b>—12.</b> 5	+13.5		╢.	
	1		-0.5 + 12.5	-18.0 -13.0			
44.07 1.01-		,,	+79.0	+13.0			
pr. 14 (Naval Observatory)	11	49	-49. 5 -42. 0	+18.0 +11.0		i i	
			-30.5	+32.5		. 1	
			-27.5 +11.5	-15.0 -21.0		1	
			+12.5	+14.5		. 1	
			+21.5 $+24.0$	+14.5 19.0		: ا	
	1		+39.5			1 3	

<sup>&</sup>lt;sup>1</sup> Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere.

Positions and areas of sun spots-Continued

Positions and arears of sun spots-Continued

	Eastern	Heliog	raphic	Aı	rea	Eastern		Heliog	iographic		Area	
Date	standard civil time	Longi- tude	Latitude	Spot	Group	Date	standard civil time	Longi- tude	Latitude	Spot	Group	
	h. m.	۰	•				h. m.		0	1.00		
Apr. 15 (Naval Observatory)	11 42	-37. 0 -28. 0	+18.0 +11.0	31		Apr. 20 (Naval Observatory)	11 49	+44.0 +49.5	-10.5 +30.0	170 15		
	ļ	-27. 0 -13. 5	-12.0 +30.0	40	154	Apr. 21 (Naval Observatory)	11 40	+57.0 $-60.0$	-13.5 $-20.0$	123	12	
		-13.5 $+25.0$	-15.0	25	216			-32.0 -22.0	+9.5 +24.5	31		
		+27.0	-11.0 $+14.5$		123			+48.0 +58.5	-10.5 -10.5	139 231		
		+34.5 +37.0	+14. 5 -19. 0	18	340			<b>∔70.0</b>	-13.5	201	. 12	
Aprl 16 (Naval Observatory)	11 41	+52.5 -21.0	$-12.5 \\ +21.0$			Apr. 22 (Mount Wilson)	18 30	-78.0 -43.5	-15.0 -20.0	55	- 2	
April 10 (1424a) Obbol 420013/2-2-2-2-	1	-17. 5 -13. 0	+13.0 -10.0		. 15			-15.0 -5.0	+8.0 +24.0	28		
	1	-10.0	+12.0		46	Apr. 23 (Naval Observatory)	11 52	+70.0 -58.5	-12.0		37	
		-5.0 -2.0	+36.5 -14.0		. 185	Apr. 23 (Navai Observatory)	11 00	-34.0	-20.0	58	1	
		+2.0 +37.0	+31.0 -23.5					-6.5 +3.0	+24.5	25	_	
		+43.0 +49.5	+12.5					+74.0 +85.0	-10.0 -10.0	154 247		
		+62.0	-16.0		62	Apr. 24 (Naval Observatory)	12 28	-46.0 -20.0	-17.0		. 10	
Apr. 17 (Naval Observatory)	- 11 42	+1.0 +11.0	+30.5	31				+16.5	+24.0		.) 10	
,		+13.0 +54.0	-14.0 $-21.0$		139 46	Apr. 25 (Naval Observatory)	11 47	-76.5 -33.0	-17.5		_ 20	
		+54.0 +63.0	+14.0				1	-7.0 +29.0		46	<u> </u>	
		+78.0	-11.5	62		Apr. 26 (Yerkes)	18 16	-59.0 -41.5	+4.0	75	i	
Apr. 18 (Naval Observatory)	11 44	-73. 8 +14. 8	-10.5		_ 370			-16.0	-16.5		. 1	
		+24.0 +28.0	14.0		139			+9.5 +19.0	-17.5	25		
		+67. 0 +79. 0	+15.0 -17.0					+45.0 +72.0	+25.0 -8.0	50 75		
Apr. 19 (Mount Wilson)	12 40	<b>-87.</b> (	) 20.0	118		Apr. 27 (Naval Observatory)	12 15	-49. 5 -32. 0	+4.5	170		
	1 .	-57.0 -50.0	-18.0	\	8			-6.0	-16.5		. 1	
	[	+27.0 +39.0						+19. 5 +56. 5	+24.5	31		
		+45.0 +68.0			205	Apr. 28 (Naval Observatory)	- 11 47	-35. 5 -17. 5			·	
4	11 40	+86.0	+13.0	) 80	)			+3.5 +9.8	-17.5			
Apr. 20 (Naval Observatory)	11 49	-73. 4 -45. 6 +34.	)   +9.5		46			+31. 8				

## AEROLOGICAL OBSERVATIONS

By L. T. SAMUELS

The month was characterized by free-air temperatures which were above normal, particularly at the higher levels. (See Table 1.) The departures were greatest from 2,000 meters to 3,000 meters at the southern stations. As might be expected under such conditions, the relative humidity was mostly below normal and the vapor-pressure departures were in general of the same sign as those for temperature.

A general excess of southerly winds occurred in the resultants for the month at all stations. (See Table 2.) The deviations from the normal were most pronounced at Royal Center and Ellendale. At the former station the northerly resultants contained a marked easterly component up to 1,250 meters instead of the more usual westerly; at Ellendale the resultant direction was southwesterly as compared to the normal northwesterly. In this connection it is interesting to find that the total precipitation at both of these stations exceeded all previous records for April since their establishment in 1918.

That a high lapse rate is not always a precursor of precipitation is shown in the kite records of Due West for April 26. A second flight was made in the afternoon at this station on account of the rapid fall in pressure. Both flights were made on the northern side of a tongue of high pressure extending inland over the South Atlantic and Gulf States. The wind direction changed but little with altitude, being westerly to west-south-

westerly at the surface and west-northwesterly above 3,000 meters. The following lapse rates prevailed during the day.

Time	M. S. L. (m.)	°C./100 m.
7:56 to 9:37 a. m	217 to 1,757	. 98 1. 10

These superadiabatic lapse rates resulted in strong convection, but owing to the low relative humidity, which for the most part was below 50 per cent, no cumulus clouds developed, the sky remaining clear throughout the day.

The heights to which the temperatures successively increased under the influence of air emanating from the rear of a slowly moving high pressure area are shown in the following table taken from the kite records of Broken Arrow for the 25th, 26th, and 27th.

M. S. L. (m.)	25th	26th	27th
	(°C.)	(°C.)	(°C.)
233	7. 5 8. 2 5. 0 2. 7 3. 3 1. 5 -0. 2	10, 3 13, 0 17, 8 14, 1 10, 5 6, 8	17. 0 17. 0 22. 6 19. 8 16. 9 13. 0 9. 0